

# Computational Assessment Of Ion Distribution On Applying Multivariate Analysis Using Geographic Information Systems

<sup>1</sup>Musini.Venkateshwarlu, <sup>\*2</sup>Suresh Merugu, <sup>3</sup>Vijayan K Asari

<sup>1</sup>Dept of Civil Engineering, CMR College of Engineering & Technology, Medchal, Hyderabad, Telangana, India  
[venkatmusini@gmail.com](mailto:venkatmusini@gmail.com)

<sup>\*2</sup>Research & Development Centre, CMR College of Engineering & Technology, Hyderabad, Telangana, India  
[msuresh@cmrcet.org](mailto:msuresh@cmrcet.org)

<sup>3</sup>Ohio Research Scholars Endowed Chair in Wide Area Surveillance, University of Dayton, Dayton, OH, USA  
[vasari1@udayton.edu](mailto:vasari1@udayton.edu)

Received: 18<sup>th</sup> November 2021, Accepted: 16<sup>th</sup> December 2021, Published: 31<sup>st</sup> December 2021

## Abstract

Study on geochemical aspects of ground water carried out by collecting samples from the existing bore well and open wells apart from surface water bodies i.e., lakes and tanks in Miryalaguda area. Electrical conductivity, pH, TDS and concentration of di and momo-valent elements controlling the parameters of drinking water carried out. The area is composite of crystalline granites of Precambrian age. The groundwater is hard water that varies from acidic to alkaline. Dissolution of fluorine into groundwater mainly by weathering of rocks is primarily involved with water pollution and water borne ailments.

**Keywords:** *Geochemical, Water Parameters, Concentration of ions, Study area*

## Introduction

Major ion geochemical study of groundwater in study area carried out to assess groundwater quality apart from fluoride concentration used as major resource for drinking in the region. Under natural conditions, it has the self-purification process, play a large role in controlling the concentrations to a certain extent. The tremendous increase in population urbanization, industrialization, agriculture, etc is the major cause of pollution [1].

Fluoride has more impact on teeth and bones rather than other elements. Fluorine content lower than 0.6 mg/lit lead to cavitation in teeth and > than 1.2 mg/lit causes fluorosis [2, 3]. Increase of concentration in drinking water is the problem across the globe. India is also facing fluorosis in 17 states including Telangana affecting 62 million people in the form of dental skeletal and/or non skeletal fluorosis [4]. Studies carried out now and earlier indicate concentration of fluoride is due to granitic gneiss rocks [5, 6]. The present work is to demarcate the area of the region in to safe zone and unsafe zones by analyze groundwater with refer to concentration of fluoride. It is done by adopting simulation methods to reduce the usage of natural instead of chemical fertilizers to reduce groundwater contamination levels and to bring the quality of water for drinking [7]. The integrated groundwater prospects maps are preferred by using different geochemical studies [8].

The area of study is between latitude from 17°5' to 17°55' and East longitude 79°30' to 79°40' form part in the survey of India toposheet No. 56 P/9 (Fig.1). Grey and pink granites are formed by magmatic origin, are subjected to high degree of weathering. This can be inferred by the field studies and presence of boulders in plain terrain. Tropical climate in led to variation in temperature, ranging from [42°C to 47°C] very hot in summer with low humidity. The origin of these rocks is due to the "parental" magma [9].

## Materials and Methods

Groundwater analysis carried out by collecting underground source and surface water bodies (Fig.2). For using standard methods major ion concentration carried out [10]. Hydrogen ion concentration and Electrical conductivity are measured. Total Dissolved Solids (TDS) are computed by multiplying the electrical conductivity by a factor (0.55 – 0.75) depending on the relative concentration of ions. Fluoride was analyzed standard method. Total alkalinity as  $\text{CaCO}_3$ , Carbonate ( $\text{CO}_3^{2-}$ ) and bicarbonate ( $\text{HCO}_3^-$ ) were estimated. Total hardness as  $\text{CaCO}_3$  and Calcium ( $\text{Ca}^{+2}$ ) were analyzed. Magnesium concentration analyzed by taking the difference between TH and  $\text{Ca}^{+2}$  values where as Sodium and Potassium content analyzed by Flame Photometer. Chloride ion was estimated by standard  $\text{AgNO}_3$  titration and the values are mentioned in (Fig 2).

## Results & Discussion

Analysis of groundwater tabulated in (Table1). Groundwater pH varies from 7.2 to 8.5 indicate it is acidic to alkaline nature. TDS varies from 236 to 1322 mg/l hardness ranges from 125 to 1690 mg/l .Hardness of these samples were correlated with standard values [11]. Indicate 75.86% water is hard water. Carbonate and bicarbonate ions in groundwater vary from 0 to 45 mg/l. and 27.1 to 225.7 mg/l respectively (Table-2). Bicarbonate content is within the limits. Sodium varies from 21.9 to 151.8 mg/lit and it is within limit of 200 mg/lit drinking water, Potassium ranges from 0.6 to mg/lit to 53.6 mg/lit indicate suitability for drinking. Calcium ranges from 20 mg/lit to 514 mg/lit with refer to prescribed limit of calcium 75 mg/lit [12].Indicate 79.37% of groundwater samples a fall under drinking water category.

Drinking water prescribed limit is 30 mg/lit as against the sample show magnesium from 1.2 mg/lit to 147.1 mg/for lit. Indicate 56.89% of the samples are as per specification. Content of Fluoride with refer to standard 1.5 mg/lit] [13] and Optimum various from 0.8 to 1.2 mg/lit. 89% of samples it is less than 0.8 mg/lit lower indicate water is suitable for drinking [14]Concentration of fluoride correlated of certain dissolved other constituent concentration. The weathering caused by alternative wet and dry conditions of semi arid climate is responsible for the leaching of fluoride from minerals in the soils and rocks [5, 15, 6, and 16]. In the contour map Fig. 3. Distribution fluoride concentration mentioned.

Acidic climatic conditions are primary factor for fluoride concentration in groundwater. Dissolution of fluoride from minerals, ion exchange [17, 18, 19, 20], evaporative concentration, led to higher concentration of which is mainly influenced by local, regional geology and hydrological conditions. The chief source of fluoride is from minerals (Fluoride, Fluorapatite, Cryolite and Pophylite) as well as fluoride replacing OH ion ferromagnesium silicates (amphiboles and micas) and soils consisting of clay minerals [21, 22, 23, 24]. The degrees for weathering and leachable fluoride in terrain is more important indicate fluoride content in water rather than more presence from the rocks [23, 25].




Fig 1: Location map of the study area




Fig 2: Sample location map




Fig 3: Fluoride Distribution map of the study area

Table 1: Correlation matrix of the study area

	pH	Ec	TH	TDS	Mg	Na	K	Cl	Ca	CO <sub>3</sub>	HCO <sub>3</sub>	NO <sub>3</sub>	SO <sub>4</sub>	F
pH	1.000													
Ec	-0.273	1.000												
TH	-0.135	<b>0.479</b>	1.000											
TDS	-0.265	<b>0.999</b>	<b>0.471</b>	1.000										
Mg	-0.127	0.323	<b>0.833</b>	0.317	1.000									
Na	-0.042	-0.003	0.064	-0.003	0.107	1.000								
K	0.127	0.299	0.051	0.321	-0.054	-0.017	1.000							
Cl	-0.151	<b>0.456</b>	0.162	<b>0.460</b>	0.081	0.015	0.050	1.000						
Ca	-0.066	0.447	<b>0.848</b>	0.439	<b>0.453</b>	0.005	0.116	0.167	1.000					
CO <sub>3</sub>	<b>0.523</b>	-0.171	-0.019	-0.168	0.072	-0.030	0.050	-0.230	-0.089	1.000				
HCO <sub>3</sub>	<b>-0.477</b>	<b>0.654</b>	0.141	<b>0.647</b>	0.062	-0.012	0.251	0.310	0.124	-0.306	1.000			
NO <sub>3</sub>	-0.103	0.038	0.253	0.036	0.249	0.008	0.003	0.056	0.188	-0.051	-0.042	1.000		
SO <sub>4</sub>	0.076	0.258	-0.055	0.257	-0.055	-0.004	0.023	0.023	-0.055	-0.034	0.389	-0.024	1.000	
F	-0.140	0.206	-0.001	0.202	0.019	0.000	0.134	-0.076	-0.022	0.028	0.133	-0.047	-0.094	1.000

**Table 2: Statistical parameters of ground water samples**

	pH	Ec	TH	TDS	Mg	Na	K	Cl	Ca	CO <sub>3</sub>	HCO <sub>3</sub>	NO <sub>3</sub>	SO <sub>4</sub>	F
<b>Mean</b>	7.9	1165	559	749	54	70	3.6	144	137	14.1	113	27	14	0.7
<b>Median</b>	7.9	1102	460	706	35	78	1.9	126	128	15.0	107	17	12	0.7
<b>Mode</b>	8.2	1888	685	1208	35	86	0.9	121	52	15.0	159	33	12	0.6
<b>Standard Deviation</b>	0.3	469	326	305	42	35	7.2	96	81	9.8	47	79	14	0.2
<b>Skewness</b>	-0.1	0.1	1	0.2	0.9	0.4	6.1	2.7	1.9	0.5	0.2	7.3	7	5.1
<b>Range</b>	1.3	1602	1565	1085	146	130	53.0	579	494	45.0	199	606	108	1.5
<b>Minimum</b>	7.2	370	125	236	1.2	22	0.6	36	20	0.0	27	2	5	0.5
<b>Maximum</b>	8.5	1972	1690	1322	147	152	53.6	614	514	45.0	226	608	113	2.0

## Conclusion

The study reveals that the groundwater of the regions is acidic to alkaline in nature. the climatic condition of the region lead to high rate A high rate of evapotranspiration and prolonged contact with weathered zone and stagnation in the led to increase in fluoride content beyond the limit of the region.

## Acknowledgements

Authors express gratitude to Ch. Gopal Reddy, Secretary, CMRGI, Principal and HOD, CMR College of Engineering & Technology Hyderabad, Head, Department of Applied Geochemistry, Osmania University, and Hyderabad for providing necessary infrastructure and encouragement.

## References

- [1].Musini Venkateshwarlu, Suresh Merugu.2020. "Fuzzy Inference System Based Assessment of Pollution Aspects in Hyderabad" Advances in Cybernetics, Cognition, and Machine Learning for Communication Technologies (pp.579-587). DOI: 10.1007/978-981-15-3125-5\_56. Online ISBN 978-981-15-3125-5.
- [2].WHO. (1993). Guide lines for Drinking Water Quality 2<sup>nd</sup> edition Vol. 1. Recommendations, Geneva.
- [3].WHO. (1996). Guidelines for Drinking Water Quality 2<sup>nd</sup> edition. Health criteria and supporting information Geneva. 231-237.
- [4].Susheela, A.K. (1999). Fluorosis management programmer in India current Sci. 77: 1250- 1256
- [5].Ramamohan Rao, N.V, Suryaprakash Rao, K and schilling, R.D. (1993). Fluorine distribution in water of Nalgonda District, Andhra Pradesh, India.Envron.Geol.21: 84-89.
- [6].Subba Rao, N. (1998). Groundwater quality in crystalline terrain of Guntur District, Andhra Pradesh, India, Visakha, sci.j.2: 51-54.

- [7].Musini Venkateshwarlu, Suresh Merugu, T. Rohini Reddy.2020. "Information System Used to Study on Effect of Groundwater Quality". *Advances in Cybernetics, Cognition, and Machine Learning for Communication Technologies* pp 531-542.[http://doi-org-443.webvpn.fjmu.edu.cn/10.1007/978-981-15-3125-5\\_53](http://doi-org-443.webvpn.fjmu.edu.cn/10.1007/978-981-15-3125-5_53).
- [8].M. Venkateshwarlu, K.Rajagopal and Y.S.Reddy (2020), Geochemical studies on the Groundwater of kistapur village, Medchal district, Telangana. *Indian J. Environmental protection*, Vol.40.No.8, PP.887 -891.
- [9].M.Venkateshwarlu, B. Prasad, A. P. Ravichandra, M.Prashanthi. "Petrography of Granitic Rocks of, Medak District, Telangana State" .*The International journal of analytical and experimental modal analysis*. Volume XII, Issue I, January/2020 Page No: 570-573.ISSN NO: 0886-9367.
- [10].American Public Health Association (APHA), *Standard Methods for the Examination of Water and Wastewater*, 6th Edition, Washington D.C., 1985.
- [11].Sawyer, C.N. and Mc Carty, P.L. (1967). *Chemistry for pamtary engineers*. Ind. D. Mc Graw hill, New York, 518 p
- [12].ISI (1983) *Indian Standard Specification for Drinking Water*. IS: 10500.
- [13].WHO. (1984). *Guidelines for Drinking Water Quality Vol 2. Health criteria and supporting information* Geneva. 100-105.
- [14].BIS (2003), *Drinking water Specification*, Bureau of Indian Standards, New Delhi IS: 10500, 11.
- [15].Wodeyar, B.K. and Srinivasan, G. (1996). Occurrence of fluoride in the groundwaters and its impact in Peddavankahalka Basin, Bellary District, Karnataka, India. *A Preliminary study current Sci* 70: 71-74.
- [16].Saxena, V.K. and Ahmad, S. (2001). Dissolution of fluoride in groundwater, a water–rock interaction study. *Environ Geol* 40: 1084-1087.
- [17].Handa, B.K. (1975). Geochemistry and genesis of fluoride containing groundwater in India, *Groundwater*13: 275-281.
- [18].American Public Health Association (APHA), *Standard Methods for the Examination of Water and Wastewater*, 6th Edition, Washington D.C., 1985.
- [19].Liu and Zhu, WH. (1991). Environmental. Characteristics of regional groundwater in relation to fluoride poisoning in North China, *Environ, Gool. Water, Sci*, 18: 3-10.
- [20].Saxena, V.K. and Ahmed, S. (2003). Inferring the chemical parameters for the dissolution of fluoride in groundwater *J.Environ.Gool*.43: 731-736.
- [21].Subba Rao, N., Krishna Rao, G. and John Devadas, D. (1998). Variation of fluoride in groundwater of crystalline terrain, *J. Environ. Hydrol*. 6: 1-5.
- [22].Agarwal, V., Vaish, A.K. And Vaish, P. (1997). Groundwater quality focus on fluoride on fluorosis in Rajasthan, *Curro, Sci* 73: 743-746.
- [23].Ramesam, V and Rajagopalan, K. (1985). Fluoride ingestion into the natural waters of hard rock areas. Peninsular India. *J.Gool.soc.Inida*.26: 125-132.
- [24].Muralidharan, D. Nair, A.P. And Satyanarayan, V. (2002). Fluoride in shallow aquifer in Ragarh Tehsil of Churu District Rajasthan; an arid Environment *Curr.Sci*.83: 699-702
- [25].Sahu, N.K. and Karim, M.A. (1989). Fluoride incidence in natural waters in Ameeli District, Gujarat.*J.Gool.soc.India*, 33, 450-456.